



# Skimming the Lunar Surface for Science: The LADEE Mission

Brian Lewis – Spacecraft Manager

# Lunar Atmosphere and Dust Environment Explorer

## Objective

- Measure Lunar Dust
- Examine the Lunar atmosphere

## Key parameters

- Launched Sept 6, 2013
- Lunar Impact April 18, 2014

## Spacecraft



- Type: Small Orbiter - Category II, Enhanced Class D
- Provider: NASA ARC and NASA GSFC

## Instruments

- Science Instruments: NMS, UVS, and LDEX
- Technology Payload: Lunar Laser Communications Demo


Launch Vehicle: Minotaur V

Launch Site: Wallops Flight Facility




# LADEE Science Objectives

- Determine **composition** of the lunar atmosphere, investigate processes controlling distribution and variability - sources, sinks, and surface interactions.
- Characterize lunar exospheric dust environment, measure spatial and temporal variability, and impacts on the lunar atmosphere.



Sample Star Tracker Image – Orbit 193





# LADEE Science Payload



- Neutral Mass Spectrometer (NMS) measures in situ ambient lunar exospheric species (eg., Ar, He, Ne)
- Ultraviolet/Visible Spectrometer (UVS) measures emissions from exospheric species and scattered light from dust
- Lunar Dust Experiment (LDEX) measures in situ dust



# LADEE Lasercomm Payload



LLCD/OM Integrated on the LADEE Spacecraft

- Achieved 622 Mbps downlink
- 20 Mbps uplink
- Achieved fully autonomous, fully optical two-way lockup.
- Streaming video!






National Aeronautics and  
Space Administration



# Flight System Overview

Star Tracker CHUs and Baffles



- Composite facesheet / Al honeycomb structure
- Single stage biprop system
- Body mounted solar arrays
- 24 A-h Li-Ion battery
- S-Band STDN compatible comm system
- Broad Reach IAU with RAD750 PPC
- Passive thermal control with heaters
- 3-axis RWA based GNC for nominal ops
- RCS thrusters for control during burns and momentum management
- 240.7 kg dry mass with 134.8 kg prop mass
- ~1090 m/s Delta-V with ~100 Nms momentum unload capacity







National Aeronautics and  
Space Administration



# Spin Balancing LADEE





# Fueling LADEE



# LADEE Launch 6 Sept 2013



© Anthony Lee

View from Big Apple



Photo: Ben Cooper



# View from Suborbital Amphibian




Photo credit: Chris Perry of the WFF Optical Systems Group.



National Aeronautics and  
Space Administration




## LADEE's Journey to the Moon



Maneuver	Date / Time (UTC)	Baseline Magnitude (m/s)	Ops Target Magnitude (m/s)	Perf. Error (m/s)	Perf. Error (%)
AM1B	11 Sep 2013 23:00	7.0	9.35	0.03	+1.0
PM1	13 Sep 2013 16:36	20.3	16.97	-0.21	-1.2
PM2	21 Sep 2013 11:53	16.0	17.49	0	0.0
TCM1	01 Oct 2013 22:00	N/A	0.88	0.04	+4.2
LOI1	06 Oct 2013 10:57	328.2	333.4	-2.9	-0.9
LOI2	09 Oct 2013 8:16	295.8	292.9	-1.8	-0.6
LOI3	13 Oct 2013 02:57	238.8	243.4	+0.8	+0.35

# Lunar Gravity and Influence on LADEE

Orbit








National Aeronautics and  
Space Administration



## LADEE Lunar Trajectory Overview







National Aeronautics and  
Space Administration



# LADEE Argon measurements

- Apollo 17 detected argon
- Not detected by LRO
- NMS detected argon and mapped its movements
- Thin layer of Argon sticks to cold lunar surface (like frost) and is released by solar warming







National Aeronautics and  
Space Administration

Ames

Discovery • Innovations • Solutions

# LADEE Argon measurements


- Localized “bump” in  $^{40}\text{Ar}$  density at  $\sim -50^\circ$  longitude
- Persists throughout dayside
- This location is special






# UVS Viewing Geometry and Data Products

Observation A





Observation B




# Characterizing the Lunar Atmosphere: First results from LADEE Sodium observations

- UVS has constructed maps of sodium in the atmosphere
- Sodium varies with lunar phase, increasing toward full moon
- Sodium increases with meteoroid showers
- Decreases in the Earth's magnetotail
- Variations suggest multiple sources that constrain processes at other airless solar system bodies



# Characterizing the Lunar Dust Environment: First results from LADEE Dust measurements

- Over 11,000 impacts recorded
- Dust cloud sustained by bombardment from interplanetary particles
- Occasional particle bursts detected
- Likely generated from impacts close to LADEE's path





This figure shows the variation in density of impacts seen by LADEE at different heights above the surface and at different times of the lunar day.

The discovery of the lunar dust exosphere opens the door to new approaches to surface composition studies and will improve our hazard estimates of large ( $> 100 \mu\text{m}$ ) dust impacts.




## High Precision in Predicted Position

- LADEE Orbit Determination team predicted the location of the spacecraft precisely enough for an LROC photo at a high velocity fly-by
  - Two spacecraft at a nearly perpendicular orbit crossing
  - Both travelling at 1.6 km/sec



# Star Tracker Images





National Aeronautics and  
Space Administration








National Aeronautics and  
Space Administration



# LADEE's Final Days






National Aeronautics and  
Space Administration




# Predicted Height Above Lunar Terrain






# And then.....







# LADEE Impact Site



Location is eastern outer rim of Sundman V crater





National Aeronautics and  
Space Administration

Ames

Discovery • Innovations • Solutions

## BioSentinel – Mission Description

- 6U autonomous nanosatellite
  - 4U microfluidic biology payload
    - Microwells with yeast
    - Tailored to detect radiation based DNA damage
    - Includes silica based LET spectrometer
  - ~2U for bus, ADCS, and propulsion
  - 14 kg mass, 23 W power
  - 18 month duration
- Identical payload developed for ISS
- Identical payload for delayed-synchronous ground control
- Identical payload for radiation exposure ground control





National Aeronautics and  
Space Administration

# Ames

Discovery • Innovations • Solutions


## Launch



Artist's rendering  
of the Space  
Launch System

- Launched as a secondary payload on **EM-1**
  - **E**xploration **M**ission 1:  
1<sup>st</sup> flight of NASA's  
Space Launch System
- Exact deployment orbit of 2° payloads still being determined
- Will likely be Earth-leading, heliocentric orbit
- Ranges of up to 70m km from Earth
- Far outside the LEOs typically occupied by CubeSats
  - ... and far outside the protective shield of Earth's magnetosphere

## Orbit



A representative orbit that  
BioSentinel might occupy





National Aeronautics and  
Space Administration



# Who am I

